

UNIVERSITY OF CALIFORNIA

San Diego

Purification and Properties of Aldehyde Dehydrogenase

from Clostridium kluyveri

A dissertation submitted in partial satisfaction of the

requirements for the degree Doctor of Philosophy

in Chemistry

by

Linda Tombras Smith.

Committee in charge:

Professor Nathan O. Kaplan, Chairman

Professor William S. Allison

Professor Robert C. Fahey

Professor Percy J. Russell

Professor Gordon H. Sato

1977

The dissertation of Linda Tombras Smith is approved, and it is acceptable in quality and form for publication on microfilm:

Robert C. Abbey

William A. Allison

Gordon Sato

Ray Russell

N. O. Kaplan

Committee Chairman

University of California, San Diego

1977

TABLE OF CONTENTS

		Page
	List of Abbreviations	viii
	List of Figures	x
	List of Tables	xiii
	Acknowledgments	xv
	Vita, Publications and Field of Study	xvi
	Abstract	xvii
I	Introduction	1
	A. Clostridia - General Description	1
	B. Metabolism of Clostridia	3
	C. Enzymes of Energy Metabolism of <u>Clostridium kluveri</u>	10
II	Materials and Methods	13
	A. Growth of <u>Clostridium kluveri</u>	13
	B. Methods of Assay	13
	1. Aldehyde Dehydrogenase Activity	14
	2. Alcohol Dehydrogenase Activity	15
	3. NADH Oxidase Activity	15
	4. Phosphotransacetylase Activity	16
	5. Determination of Protein Concentration	16
	C. Preparation and the Determination of the Concentration of Pantetheine	17
	D. Synthesis and Determination of the Concentration of Acetyl Coenzyme A	18
	E. Affinity Chromatography	19
	1. Coupling of Amines to Sepharose	19
	2. Amide Bond Synthesis	20
	3. Preparation of a Sepharose-Bound Acetaldehyde Derivative	20

	Page
4. Preparation of Sepharose-Bound Desulfo-Coenzyme A	21
5. Binding of Aldehyde Dehydrogenase to Sepharose-Bound Ligands	22
F. Purification Procedure for Aldehyde Dehydrogenase	23
G. Polyacrylamide Gel Electrophoresis	26
H. Active Enzyme Centrifugation	28
I. Stokes Radius Determination	29
J. Thiol Specificity	30
1. Charcoal Treatment	30
2. Alkaline Phosphatase Treatment	31
3. Pyrophosphatase Treatment	32
4. Product Identification	32
K. Rapid Kinetic Measurements	33
L. Methods for Kinetic Analysis	33
1. General Methods	34
2. Determination of the Concentration of Substrates	36
M. Purification Procedure for Phosphotransacetylase	36
N. Phosphotransacetylase Activity Stain for Polyacrylamide Gels	37
III Results and Discussion	39
A. Syntheses	39
1. Preparation of Pantetheine	39
2. Synthesis of Acetyl Coenzyme A	40
3. Preparation of Sepharose-Bound Desulfo-Coenzyme A	43
B. Purification of Aldehyde Dehydrogenase	46
1. Extraction of the Enzyme	46
2. Precipitation of Activity	49
3. Desalting by Gel Filtration	52

	Page
4. DEAE-Cellulose Chromatography	52
5. Concentration Step	57
6. Calcium Phosphate Gel Adsorption	59
7. Gel Filtration	59
8. Polyacrylamide Gel Electrophoresis of Aldehyde Dehydrogenase	62
9. Discussion of the Purification Scheme	62
C. Binding of Aldehyde Dehydrogenase to Sephacrose-Bound Ligands	69
D. Stabilization of Aldehyde Dehydrogenase	77
E. Molecular Weight Determination	81
1. Determination of Sedimentation Coefficient	81
2. Determination of Stokes Radius	86
3. Calculation of Molecular Weight and Frictional Ratio	87
4. Summary of Ultracentrifugation and Gel Filtration Studies	91
F. Aldehyde Specificity	93
G. Pyridine Nucleotide Specificity	94
H. Pyridine Nucleotide Stereospecificity	97
I. Thiol Specificity	98
1. Utilization of Various Thiols	98
2. Spectral Studies and Treatment with Charcoal	100
3. Reaction in the Presence of Phosphotransacetylase	104
4. Alkaline Phosphatase Treatment	105
5. Pyrophosphatase Treatment	105
6. Thioester Formation	105
7. General Discussion	111
J. Rapid Kinetic Measurements	113
K. Kinetic Analysis	116
1. Initial Velocity Studies	116
2. Product Inhibition	139
3. Reverse Reaction	151
4. Substrate Inhibition	155
5. Discussion of Kinetic Analysis	161

	Page
L. Purification of Phosphotransacetylase	167
1. Results and Discussion of Purification Procedure	167
2. Polyacrylamide Gel Electrophoresis of Phosphotransacetylase	172
References	179
Appendix	187

ACKNOWLEDGMENTS

I would like to thank Dr. Nathan O. Kaplan for his scientific guidance, support, and encouragement. Dr. G. D. Novelli is gratefully acknowledged for providing the Clostridium kluyveri cells. Thanks also to Dr. J. Everse for assistance with the ultracentrifugation and stopped-flow studies. Drs. G. M. Smith and C. Y. Lee ran and assisted in the interpretation of the nuclear magnetic resonance spectra. Thanks to Dr. Y. Levin for providing his expertise in the field of affinity chromatography. Drs. L. J. Arnold and K. S. You are acknowledged for their helpful advice.

VITA

March 1, 1950, Born, Oakland, California

1967-1969 Diablo Valley College, Pleasant Hill, California
1971- B.A., University of California, Berkeley
1972-1975 Teaching Assistant, Department of Chemistry,
 University of California, San Diego
1972-1977 Research Assistant, Department of Chemistry,
 University of California, San Diego
1977- Doctor of Philosophy

PUBLICATIONS

"Purification and Characterization of Aldehyde Dehydrogenase from Clostridium kluveri." Fed. Proc., 35: 1750 (1976).

FIELD OF STUDY

Major Field: Biochemistry

Studies in Enzymology

Professor Nathan O. Kaplan

ABSTRACT OF THE DISSERTATION

Purification and Properties of Aldehyde Dehydrogenase

from Clostridium kluveri

by

Linda Tombras Smith

Doctor of Philosophy in Chemistry

University of California, San Diego, 1977

Professor Nathan O. Kaplan, Chairman

Coenzyme A-linked aldehyde dehydrogenase from Clostridium kluveri catalyzes the oxidation of acetaldehyde in the presence of CoA and oxidized pyridine nucleotide to form acetyl CoA and reduced pyridine nucleotide. The enzyme was purified approximately 90 fold over crude extracts to a specific activity of 50 units/mg protein. From polyacrylamide gel electrophoresis, the enzyme preparation was estimated to be 35% pure. Although the dehydrogenase showed some affinity toward substrate and hydrophobic ligands bound to Sepharose, none of the gels tested were found to be useful for purification purposes. The enzyme was partially stabilized by thiols at low temperatures.

From active enzyme centrifugation studies, aldehyde dehydrogenase was found to have a sedimentation coefficient of $S_{20,w} = 7.4$. From gel filtration studies, the Stokes radius of the enzyme in the presence of substrates was found to be 9.5 nm and in the absence of substrates was 11.0 nm. Using the values found for the sedimentation coefficient and the Stokes radius, the molecular weight of the enzyme in the presence of substrates was calculated to be 290,000 daltons. The frictional ratio was 2.2.

Aldehyde dehydrogenase can utilize both NAD^+ and NADP^+ to catalyze the formation of acetyl CoA. The rate obtained with NADP^+ was about 30% of that obtained with NAD^+ . To a limited extent, the dehydrogenase may utilize thiols other than CoA as acetyl acceptors. A number of methods were employed in order to exclude the possibility that these thiols act merely by recycling trace amounts of CoA that might be in the enzyme preparation via nonenzymatic transacetylation.

Initial velocity and product inhibition studies were conducted in order to elucidate the order of substrate binding and product release. A Bi Uni Uni Uni Ping Pong kinetic mechanism was proposed in which NAD^+ binds to the enzyme first, acetaldehyde second, and NADH is released. Then CoA binds and acetyl CoA is released. At K_M levels of NAD^+ and CoA, substrate inhibition by CoA was observed. It was suggested that CoA binds to free enzyme to form a dead-end E·CoA inhibition complex.

Phosphotransacetylase from Clostridium kluyveri was purified more than 140 fold over crude extracts, by a combination of ammonium sulfate fractionation and affinity chromatography with C⁸-(aminohexyl)-amino-desulfo-CoA bound to Sepharose. The purified protein was judged to be over 70% pure by polyacrylamide gel electrophoresis.